

Global analysis of depletion and recovery of seabed biota after bottom trawling disturbance

Jan Geert Hiddink^{a,1}, Simon Jennings^{b,c,2}, Marija Sciberras^a, Claire L. Szostek^a, Kathryn M. Hughes^a, Nick Ellis^d, Adriaan D. Rijnsdorp^{e,1}, Robert A. McConnaughey^f, Tessa Mazor^d, Ray Hilborn^g, Jeremy S. Collie^h, C. Roland Pitcher^d, Ricardo O. Amorosoⁱ, Ana M. Parma^j, Petri Suuronen^j, and Michel J. Kaiser^a

^aSchool of Ocean Sciences, Bangor University, Anglesey LL59 5AB, United Kingdom; ^bCentre for the Environment, Fisheries and Aquaculture Science, Lowestoft, Suffolk NR33 0HT, United Kingdom; ^cSchool of Environmental Sciences, University of East Anglia, Norwich Research Park NR4 6TPF, United Kingdom; ^dEcoSciences Precinct 41, Commonwealth Scientific and Industrial Research Organization Oceans & Atmosphere, Brisbane, QLD 4001, Australia; ^eInstitute for Marine Resources and Ecosystem Studies, Wageningen UR, 1700 AB IJmuiden, The Netherlands; ^fResource Assessment and Conservation Engineering Division, Alaska Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, Seattle, WA 98115; ^gSchool of Aquatic and Fishery Sciences, University of Washington, Seattle, WA 98195; ^hGraduate School of Oceanography, University of Rhode Island, Narragansett, RI 02882; ⁱCentro Nacional Patagónico, National Scientific and Technical Research Council (CONICET), 9120 Puerto Madryn, Argentina; and ^jFisheries and Aquaculture Department, Food and Agriculture Organisation of the United Nations, 00153 Rome, Italy

Edited by Nils C. Stenseth, University of Oslo, Oslo, Norway, and approved June 12, 2017 (received for review November 15, 2016)

Bottom trawling is the most widespread human activity affecting seabed habitats. Here, we collate all available data for experimental and comparative studies of trawling impacts on whole communities of seabed macroinvertebrates on sedimentary habitats and develop widely applicable methods to estimate depletion and recovery rates of biota after trawling. Depletion of biota and trawl penetration into the seabed are highly correlated. Otter trawls caused the least depletion, removing 6% of biota per pass and penetrating the seabed on average down to 2.4 cm, whereas hydraulic dredges caused the most depletion, removing 41% of biota and penetrating the seabed on average 16.1 cm. Median recovery times posttrawling (from 50 to 95% of unimpacted biomass) ranged between 1.9 and 6.4 y. By accounting for the effects of penetration depth, environmental variation, and uncertainty, the models explained much of the variability of depletion and recovery estimates from single studies. Coupled with large-scale, high-resolution maps of trawling frequency and habitat, our estimates of depletion and recovery rates enable the assessment of trawling impacts on unprecedented spatial scales.

logistic recovery model | systematic review | metaanalysis | impacts | trawling

Fisheries using bottom trawls are the most widespread source of anthropogenic physical disturbance to global seabed habitats (1, 2). Almost one-quarter of global seafood landings from 2011 to 2013 were caught by bottom trawls (3). Development of fisheries, conservation, and ecosystem-based management strategies requires assessments of the distribution and impact of bottom trawling and the relative status of benthic biota and habitats. There are many drivers for such assessments, including (i) policy commitments to an ecosystem approach to fisheries, (ii) requirements to take account of trawling impacts in fisheries and environmental management plans, (iii) demands from certification bodies to assess fisheries' environmental impacts, and (iv) the need to evaluate the effects of alternate management measures to meet conservation and management objectives (4–6). These assessments are used to assess the sustainability of bottom trawl fisheries, formulate priorities for habitat protection, and ultimately, achieve a balance between fisheries production and environmental protection. The distribution of bottom trawling is increasingly well-characterized by vessel tracking and other monitoring systems (7), but impacts depend on the magnitude of trawling-induced mortality and recovery rates of biota, for which the current evidence base is incomplete, dispersed, and often contested (4, 8).

Bottom trawls [here defined as any towed bottom-fishing gear, including otter trawls (OTs), beam trawls (BTs), towed (scallop) dredges (TDs), and hydraulic dredges (HDs)] are used to catch fish, crustaceans, and bivalves living in, on, or above the seabed (9). Bottom trawling resuspends sediments (10, 11); reduces topographic complexity and biogenic structures (12–14); reduces faunal

biomass, numbers, and diversity (15, 16); selects for communities dominated by fauna with faster life histories (17); and produces energy subsidies in the form of carrion (18). These effects lead to changes in community production, trophic structure, and function (19, 20). Given the patchy and dynamic distribution of bottom fishing (21), fished seabeds comprise a mosaic of undisturbed, recently impacted, and recovering benthic communities and habitats (22). The state of each patch within this mosaic depends on the history and frequency of past trawling impacts and the recovery rates of the biota present (23).

Recovery rates after trawling depend on recruitment of new individuals, growth of surviving biota, and active immigration from adjacent habitat. Most existing estimates of recovery rates come from experimental studies, with changes in abundance recorded before and after experimental trawling (15, 16). Although these experiments provide reliable estimates of immediate mortality, their small scale is likely to underestimate recovery time, in particular for mobile fauna. This underestimation is because immigration makes a greater contribution to recovery when biota are relatively more abundant around

Significance

Bottom trawling is the most widespread source of physical disturbance to the world's seabed. Predictions of trawling impacts are needed to underpin risk assessment, and they are relevant for the fishing industry, conservation, management, and certification bodies. We estimate depletion and recovery of seabed biota after trawling by fitting models to data from a global data compilation. Trawl gears removed 6–41% of faunal biomass per pass, and recovery times posttrawling were 1.9–6.4 y depending on fisheries and environmental context. These results allow the estimation of trawling impacts on unprecedented spatial scales and for data poor fisheries and enable an objective analysis of tradeoffs between harvesting fish and the wider ecosystem effects of such activities.

Author contributions: J.G.H., S.J., R.H., and M.J.K. designed research; J.G.H., S.J., M.S., C.L.S., K.M.H., N.E., R.A.M., R.H., J.S.C., C.R.P., R.O.A., A.M.P., P.S., and M.J.K. performed research; J.G.H. analyzed data; and J.G.H., S.J., M.S., C.L.S., N.E., A.D.R., R.A.M., T.M., R.H., J.S.C., C.R.P., R.O.A., A.M.P., P.S., and M.J.K. wrote the paper.

Conflict of interest statement: The study was funded by a variety of organizations that include nongovernmental organizations that promote conservation and sustainable use and fish producers.

This article is a PNAS Direct Submission.

Freely available online through the PNAS open access option.

¹To whom correspondence may be addressed. Email: j.hiddink@bangor.ac.uk or Adriaan. Rijnsdorp@wur.nl.

²Present address: International Council for the Exploration of the Sea, 1553 Copenhagen V, Denmark.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1618858114/-DCSupplemental.

Hiddink et al.




Fig. 2. The relationship between the penetration depth P and depletion d of macrofaunal community biomass and numbers caused by a single trawl pass for different trawl gears (means \pm SD).

Table S5), but the model estimates for gravel are nevertheless shown in Table 1 for community biomass to allow comparison with the significant effects of gravel found for community numbers (Table 1, community numbers and *SI Appendix, Table S5*). Mean community r (estimated using *SI Appendix, Eqs. S4.1 and S4.2* from d and b) increased with trawling frequency from 0.82 y^{-1} when there was no trawling (5–95% uncertainty intervals = $0.42\text{--}1.53$) to 1.73

$(0.89\text{--}3.23) \text{ y}^{-1}$ when the trawling frequency was 10 y^{-1} (using the mean estimated d across gears OT, BT, and TD; $d = 0.13$) (SI Appendix, Fig. S1 and Table S6). The increase in r , which results from changes in community composition to favor biota with faster life histories, is, therefore, relatively slight across ranges of trawling frequencies that dominate those on real fishing grounds (e.g., $0\text{--}1 \text{ y}^{-1}$) (7, 31, 32). The r estimate of 0.82 y^{-1} enables estimates of median time to recovery (T) to 0.95K for a range of levels of depletion (Fig. 3B). For example, if the fraction depleted $D = 0.5K$, then recovery time is 3.6 y (5–95% uncertainty intervals = 1.9–6.4 y).

The effect of trawling on community numbers, estimated from the comparative studies, increased significantly with the gravel content of the sediment (Fig. 3C, Table 1, community numbers, and *SI Appendix, Table S5*), and this effect persisted when examined among gears. The reductions in benthic community numbers for each unit increase in trawling frequency were 3.1% at 0% gravel content (typical for BT studies), 5.5% at 1% gravel content (typical for OT studies), and 72% at 45% gravel content (typical for TD studies). The estimates of r for community abundance range from 0.18 y^{-1} for TD on 45% gravel to 4.47 y^{-1} for BT on 0% gravel, with high uncertainty. These r estimates result in a median recovery time T from $0.5K$ to $0.95K$ of $0.7\text{--}16.6 \text{ y}$ (Fig. 3D). Other than gravel content, the inclusion of the ratio of d over primary production also resulted in reduced Akaike information criterion (AIC) compared with the model with no additional explanatory variables, with the effect of trawling on numbers increasing with d and decreasing at higher levels of primary production (Table 1, community numbers and *SI Appendix, Table S5*).

Discussion

This study is an attempt to quantify the impacts of bottom trawling and recovery of seabed biota by synthesizing data from trawling studies after a systematic review of the available evidence base. We developed a method to derive the recovery rates of benthic macro-faunal invertebrate communities from trawling by combining results from experimental and comparative studies and provide estimates of depletion and recovery, including a quantification of uncertainty




Fig. 3. The relationship between trawling frequency and total community (A) biomass and (C) numbers. The thicker lines are the fixed effects, and gray lines are the random effects of the individual studies (not all visible, because many studies had small ranges and low trawling frequencies). Recovery time to 0.95K for depleted total community (B) biomass and (D) numerical abundance as a function of estimated r and initial depletion D . In A and B, lines are the median estimate based on the mean d across all gears. In C and D, lines are the median estimates for three different gear types based on the mean gravel content in the areas where studies using these gear types were carried out. The shaded areas indicate the 5–95% uncertainty intervals for estimates.

Table 1. Linear mixed model (SI Appendix, Eq. S3.1) fits for the analysis of data from comparative studies of changes in biomass and numbers

Model	Slope (<i>b</i>)	SE	df	<i>t</i> Value	<i>P</i> value	AIC
Community biomass						
TF	−0.07522	0.0158	503	−4.732	<0.0001	566.9
TF	−0.07142	0.0172	502	−4.148	<0.0001	568.4
TF: gravel	−0.00067	0.0010	502	−0.648	0.5168	
TF	−0.08623	0.0325	502	−2.653	0.0082	568.8
TF: <i>d</i> /PP	125.6879	373.7966	502	0.336	0.7368	
Community numbers						
TF	−0.21185	0.1342	141	−1.577	0.1169	89.5
TF	−0.01451	0.0942	140	−0.153	0.8778	81.1
TF: gravel content	−0.01206	0.0035	140	−3.377	0.0009	
TF	0.25300	0.2145	140	1.048	0.2964	86.1
TF: <i>d</i> /PP	−6,892.96900	2,676.5453	140	−2.575	0.0111	

For community biomass, the model with the lowest AIC included no explanatory variables other than trawling frequency, but for community abundance, both gravel content and *d*/PP improved the AIC in relation to a model without other explanatory variables. Results for these variables are given under community biomass for comparative purposes. *d*, Depletion estimate from experimental studies (fraction per trawl pass) (SI Appendix, Table S4); gravel, sediment composition in percentage by weight; PP, primary production (milligrams C meter^{−2} day^{−1}); TF, trawling frequency.

based on all available data. The method for estimating the recovery rate from comparative studies is unique. Given that realistic and robust *r* estimates have been largely unavailable previously, this work is critically important. Recovery rates were estimated from changes in the biomass and numbers of biota across fishing grounds, and therefore, estimates are likely applicable to trawled shelf seas in general (at least in temperate waters where most of the studies were carried out). Our estimates of depletion and recovery enable the parameterization of models to predict the state of the benthic biota as a function of trawling frequency and levels of primary production and percentage gravel (28). Coupled with the emergence of large-scale estimates of trawling frequency (7), these models will support assessment of trawling impacts on unprecedented spatial scales, because our approach provides a quantitative estimate of status with minimal data requirements (28). The method is widely applicable, because it requires relatively few data inputs and could be applied worldwide, including for fisheries where trawl impacts remain unassessed. The *r* and *d* values that we estimate here with a broad geographic basis are based on the full body of available evidence and therefore, the most robust estimates available. The generality of our approach means that the outputs of assessments are accurate when averaging over larger scales but that biases may exist when used for local assessments. These results have global policy relevance for conservation and food security policy development, because they enable an objective analysis of the efficacy of different methods of harvesting food from the ocean to be considered in the light of the wider ecosystem effects of such activities on the marine environment. The results enable managers to understand the variable resilience of benthic systems to trawl fisheries and set limits of fishing accordingly.

Most continental shelves consist of relatively small intensively trawled areas, where the trawling frequency is in the range of 1–10 y^{−1}, and extensive infrequently trawled areas, where the trawling frequency is <1 y^{−1} and predominantly <0.25 y^{−1} (7). Our results show that trawling frequencies of 1 y^{−1} cause average declines of 15.5% in the biomass of benthic biota. Communities on gravel may be more sensitive to trawling, because they, on average, have a larger proportion of larger, long-lived, and sessile epifauna (33) that are particularly sensitive to trawling (34). Effects were

greater for gears that kill a larger fraction of the biota (larger *d*), because they penetrate the sediment more deeply and weaker in areas of higher primary production, where higher food supply to the benthos may result in a higher recovery rate.

The ranking of different fishing gears with respect to their magnitude of impact reported here is similar to the ranking in previous metaanalyses of small-scale experimental studies (15, 16), although our estimates of *d* are smaller, probably because we adjusted for the number of trawl passes, whereas previous analyses did not. The use of depletion to primary production ratio as a proxy for community resilience to trawling has the advantages of being easily understandable and easy to estimate for new areas and fisheries. The ratio of depletion over primary production might support rapid preliminary large-scale risk assessments of potential trawling impacts on community abundance to guide more region-specific studies. The close relationship between penetration depth and depletion can be used to estimate depletion resulting from the pass of a given trawl gear when no direct depletion estimate is available. Accurate estimates of penetration depth are much easier and cheaper to obtain than estimates of depletion, would support preliminary impact assessments by gear type, and can even be generated using numerical models (11).

Our analyses did not identify any variables other than trawling frequency that affected community biomass. This finding is surprising given the contrasting results for numbers and that some comparative studies and past metaanalyses of experimental studies have shown interaction effects between gear type and habitat type (16, 29). The relatively small number of studies included in the biomass analysis and the high variability associated with benthic sampling, which cannot be fully controlled in a metaanalysis, may have contributed to this discrepancy. Our results for biomass imply that a single estimate of recovery rate *r* is appropriate when assessing impacts on the different habitat types studied here. They also suggest that differences in time to recovery and expected biomass (*B*/*K*) will be driven primarily by gear type (and hence, *d*) and trawling frequency (*F*).

Our estimates of biomass recovery times are similar to empirical measurements of recovery taken in three areas where commercial trawling was stopped (4–5 y) (24) but longer than estimates from small-scale experimental studies, which are on the order of 25–500 d (15, 16). The scale dependency of recovery times has important implications for management, because recovery will be faster when trawled areas are closer to less impacted areas from which individuals can recruit or migrate (as also shown in ref. 22). We found that biomass recovery rates were slower and that recovery times were longer than those for numbers. This result is expected based on the population dynamics of seabed biota. Recovery in numbers is driven more strongly by recruitment than recovery of biomass, which is driven by increases in the size and age structure of the population through growth of individuals. We recommend the use of recovery rates for community biomass when modeling trawl impacts and their consequences. This approach will give due weight to recovery of body size and age structure as well as numbers and take account of energy flow through food webs and other ecosystem processes that are linked closely to biomass. Recovery times as estimated from the logistic model nevertheless do not imply that the communities will recover over these times to the species, size, and age composition that existed before trawling, but they do imply the recovery of total biomass or numbers and related cross-species ecosystem processes, such as aggregate secondary production.

Uncertainties around mean/median estimates of penetration depth, recovery, and depletion were high, despite the careful screening of included data (which also decreased the sample size and potentially, power to detect effects) (30). However, our approach allows us to address directly some aspects of uncertainty, and the broad distribution of resulting depletion and recovery estimates show that large site-specific differences in the response of seabed communities to trawling are expected. The advantage of characterizing uncertainty is that it can be propagated in future

risk and impact analyses. Given the unexplained variance in r , percentiles from the distribution of plausible values might be selected to reflect the degree of risk aversion in the management system. The extent of risk aversion is a nonscientific decision (although it would be informed by science) that would likely be made by managers and other stakeholders. Risk aversion would likely depend on the perceived value of a habitat type. A risk-averse approach might adopt a value of r from a lower percentile of the distribution (e.g., the 10 or 25%) rather than the median (*SI Appendix, Table S6* shows a selection of values).

Our use of comparative studies provides improved estimates of recovery compared with those from previous small-scale experiments studies, because they are based on larger-scale measurements from fishing grounds. Comparative studies may, however, be affected by “shifting baselines” (35), where historical trawling has removed the most sensitive organisms and only resilient organisms remain. Because trawling selects for species with faster life histories that are more resilient, recovery time will increase with trawling frequency. Our finding that mean community r increases with F conforms with previous observations of shifts toward species with faster life histories in disturbed communities (36). This effect is apparent across a range of plausible trawling frequencies from >0 to 10 y^{-1} but would be small for the great proportion of most fishing grounds, where swept area ratio is less than 1 y^{-1} (7). Although this shift means that previously trawled communities may be more resilient to additional trawling, it does not mean that they will recover any faster to the original pre-trawling state. For this reason, we used the r estimate of untrawled communities for estimating recovery times. Selective effects linked to trawling history are likely to be strongest for long-lived sessile epifauna that build biogenic reefs, such as sponges and corals. The estimates of r and T presented here are applicable to invertebrate communities living in sedimentary habitats but not biogenic habitats, because no studies of trawling impacts on biogenic habitats met the rigorous selection criteria imposed by the systematic review.

In summary, we apply widely applicable methods to estimate depletion and recovery rates of benthic invertebrate communities after trawling. By accounting for the effects of gear type and penetration, environmental variation, and uncertainty, our analysis explained much of the variability of depletion and recovery estimates from single studies. Coupled with large-scale, high-resolution maps of trawling frequency and habitat, our estimates of depletion and recovery rates will enable analysis of trawling impacts on unprecedented spatial scales to inform best practices to achieve sustainable fishing and will be of use to policymakers, conservation planners, and fisheries managers for risk assessment and the evaluation of management strategies.

Methods

We present analyses for whole-community biomass and numbers of benthic invertebrates. Changes in the abundance of seabed biota after trawling depend on the mortality caused by each pass of a trawl and the rate of recovery of the biota between trawl passes. We estimated the immediate depletion of biota (d) caused by a trawl pass from a metaanalysis of experimental studies of trawling impacts. We estimated the recovery rates (r) of biota from a metaanalysis of comparative studies of trawling impacts. The analyses were structured to assess the effects of gear type, penetration depth, and environmental variables (e.g., depth and sediment composition) on depletion and recovery.

Depletion. Depletion was estimated using data collated from experimental studies of trawling impacts identified using systematic review methodology. A comprehensive literature search of journal papers, book chapters, and grey literature reports was carried out. Details of literature search terms, databases, and study inclusion criteria are provided in the systematic review protocol by Hughes et al. (30). All included studies quantified the immediate mortality of biota after one or multiple trawling events. Each identified study had to pass quality assurance criteria before data from the study were included in the collated dataset.

We classified gear types as OTs, BTs, TDs, or HDs (*SI Appendix*). The reduction in abundance of biota resulting from one pass of a trawling gear depends on the characteristics and operation mode of the gear. Different gears are designed to have different levels of seabed contact or penetration depending on the

target species and seabed type, and these factors will influence mortality (37). Consequently, we assessed the relationship between mortality and penetration depth of the gear. Some of these studies were conducted in previously trawled areas with a lowered abundance of biota, but because we are estimating the fraction of organisms removed rather than the absolute amount, we expect that this will have had little effect on our estimates of d . Depletion d was estimated using a generalized linear mixed model implemented in the package nlme in R (38, 39), with the log of the ratio of the biomass or abundance in trawled over untrawled areas ($\ln RR$) as the response variable, \log_2 (time t in days since trawling) and gear type as fixed factors, and the study as a random effect assuming a Gaussian error distribution. We weighted $\ln RR$ values by the inverse of their variance, which is normal practice in metaanalyses. We estimated d as the intercept for the different gears at $t = 0$.

Predicted penetration depth of each gear type into the seabed was estimated from values in the literature by averaging the reported penetration depths of the individual components of the gear (e.g., doors, sweeps, and bridles of an OT) weighted by the width of these components (details are in *SI Appendix*).

Recovery. Recovery rates were estimated using data collated from comparative studies of trawling impacts. All included studies sampled the biomass or numbers of whole communities of benthic invertebrates at two or more sites subject to different trawling intensities on commercial fishing grounds. Contributing studies were identified following the same procedure as for experimental studies (*SI Appendix*). In the analyses of the comparative studies, we assume that both K and observed gradients of trawling effort were unrelated to other environmental drivers and that the observed state of the biota is in equilibrium with the reported trawling effort. Gradients in trawling effort may be driven by regulation and seabed obstructions but are also observed in areas of homogenous habitat (29). Spatial patterns of trawling effort are also shown to be relatively stable over time in the few fisheries where high-resolution time series have been analyzed (40). K could vary across the trawl grounds because of environmental variations, and this source of variation will increase the uncertainty around relationships between B and F .

In the comparative studies, conversions between units of abundance were not always possible (e.g., biomass per unit sediment volume could not be converted to biomass per unit sediment area given sampling gears with different but unknown efficiencies), and therefore, absolute B or K could not be estimated. We normalized the data by expressing relative biomass or numbers as the B/K ratio and used a log-linear approximation for the relationship between community B/K and F :

$$\log_{10}\left(\frac{B}{K}\right) \sim bF, \quad [2]$$

where b is the slope of the relationship (derivation taking account of the log-linear relationship between B/K and F and the distribution of trawling is in *SI Appendix*). After fitting a linear relationship to $\log_{10}B$ vs. F for each comparative study, K was estimated as the $10^{\text{intercept}}$ of this relationship.

The data collated from comparative studies were initially used to estimate relative changes in abundance (B/K) as a function of trawling frequency F . This approach differs from the aforementioned analyses of depletion, because the change in abundance with trawling is a response to both depletion (per trawl pass) and recovery. Because $b = d/r$ (Eq. 1), after d is estimated from experimental data, recovery rate r can be estimated from the slope b of Eq. 2 after taking account of the log-linear nature of this relationship, which implies that r increases with F . To propagate uncertainty in the estimates of b and d into the estimate of r , we sampled the distributions of b and d estimates to derive the distribution of r (*SI Appendix*). Time to recovery from a given level of depletion D to a defined proportion ϕ of K at which recovery is deemed to have occurred (e.g., 0.95) was derived from the approach of Lambert et al. (22) (*SI Appendix*). When reporting recovery times, we report recovery from $0.5K$ to $0.95K$.

Variables That Determine the Effect of Trawling in Comparative Studies. The effect of trawling on seabed biota in comparative studies could be influenced by different variables. Thus, we evaluated the explanatory power of several potential factors by including them as covariates in a linear mixed model (39) based on Eq. 2 and selecting the most parsimonious model using AIC. According to Eq. 2 the community response to trawling in \log_{10} scale is approximately proportional to F , with slope a function of the ratio of d/r . The fixed part of the mixed models was, therefore,

$$\log_{10}(\text{Response}) \sim \text{Trawling frequency} + \text{Trawling frequency} \times \text{other variables},$$

where the response variable is community biomass or numbers and the “other variables” can be covariates for d , r , or their ratio (4). The intercept

was removed, because $\log_{10}(B/K)$ with no impact = 0. We modeled "study" as a random effect, allowing the slope to vary per study. This approach accounted for the nonindependence of observations within a study. We checked the assumptions of the linear mixed model by visual inspection of the normalized residuals (38).

We expected that factors that lead to a higher d would strengthen the effect of trawling (e.g., higher penetration depth), whereas factors that lead to a higher r by affecting growth rates of individuals and populations (higher flow of energy to the seabed because of a higher production, shallower depth, or higher temperature) would weaken the effect. The closely related penetration depth P (continuous) and gear type (categorical) were examined as covariates for d . The following covariates for r were examined: primary production estimated from the vertically generalized productivity model (milligrams C meter⁻² day⁻¹) (41) and particulate organic carbon flux to depth (grams C_{org} meter⁻² year⁻¹) (42) as proxies for energy availability, mean sea bottom temperature calculated from monthly mean bottom temperature for 2009–2011 provided in MyOcean Product (GLOBAL-REANALYSIS-PHYS-001–009), depth (from GEBCO if not reported in the original study), habitat type, and sediment composition (gravel, sand, and mud content). Habitat types were classified as biogenic habitats, gravel, sand, muddy sand/sandy mud, and mud. Sediment gravel, sand, and mud content were extracted from the source studies by converting the sediment description

to the Folk classification (43) and then converting the Folk classification to percentages based on the means in each category. In addition to analyses using covariates of d or r , we also conducted analyses using covariates of the d/r ratio; here, the d/r ratio was approximated as the ratio of d or P to the continuous r covariates. The effect of trawling is expected to increase with water depth owing to the lower levels of natural disturbance in deeper water and the corresponding increase in the relative abundance of individuals with slower life histories (low r), and therefore, $d \times$ depth was examined as a covariate for d/r , with depth expressed as a negative number.

ACKNOWLEDGMENTS. The study was funded by the David and Lucile Packard Foundation; the Walton Family Foundation; The Alaska Seafood Cooperative; American Seafoods Group; Blumar Seafoods Denmark; Clearwater Seafoods; Espersen Group; Glacier Fish Company LLC; Gortons Inc.; Independent Fisheries Limited N.Z.; Nippon Suisan (USA), Inc.; Pacific Andes International Holdings, Ltd.; Pesca Chile, S.A.; San Arawa, S.A.; Sanford Ltd. N.Z.; Sealord Group Ltd. N.Z.; South African Trawling Association; and Trident Seafoods. Additional funding was provided by United Kingdom Department of Environment, Food and Rural Affairs Project MF1225 (to S.J.), European Union Project BENTHIS EU-FP7 312088, The International Council for the Exploration of the Sea Science Fund, and the Food and Agriculture Organisation of the United Nations.

- Halpern BS, et al. (2008) A global map of human impact on marine ecosystems. *Science* 319:948–952.
- Foden J, Rogers SI, Jones AP (2011) Human pressures on UK seabed habitats: A cumulative impact assessment. *Mar Ecol Prog Ser* 428:33–47.
- FAO (2016) *The State of World Fisheries and Aquaculture 2016* (FAO, Rome).
- Rice J (2005) Challenges, objectives, and sustainability: Benthic community, habitats and management decision-making. *Am Fish Soc Symp* 41:41–58.
- Rice J, et al. (2012) Indicators for Sea-floor Integrity under the European Marine Strategy Framework Directive. *Ecol Indic* 12:174–184.
- Rijnsdorp AD, et al. (2016) Towards a framework for the quantitative assessment of trawling impact on the seabed and benthic ecosystem. *ICES J Mar Sci* 73:i127–i138.
- Eigaard OR, et al. (2016) The footprint of bottom trawling in European waters: Distribution, intensity and seabed integrity. *ICES J Mar Sci* 74:847–865.
- Gray JS, Dayton P, Thrush S, Kaiser MJ (2006) On effects of trawling, benthos and sampling design. *Mar Pollut Bull* 52:840–843.
- Sainsbury JC (1986) *Commercial Fishing Methods* (Fishing News Books, Oxford).
- Churchill JH (1989) The effect of commercial trawling on sediment resuspension and transport over the Middle Atlantic Bight continental shelf. *Cont Shelf Res* 9:841–864.
- O'Neill F, Ivanović A (2016) The physical impact of towed demersal fishing gears on soft sediments. *ICES J Mar Sci* 73:i5–i14.
- Peterson CH, Summerson HC, Fegley SR (1987) Ecological consequences of mechanical harvesting of clams. *Fish Bull* 85:281–298.
- Collie JS, Escanero GA, Valentine PC (1997) Effects of bottom fishing on the benthic megafauna of Georges bank. *Mar Ecol Prog Ser* 155:159–172.
- Collie JS, Hermesen JM, Valentine PC, Almeida FP (2005) Effects of fishing on gravel habitats: Assessment and recovery of benthic megafauna on Georges Bank. *Benthic Habitats and the Effects of Fishing*, eds Barnes P, Thomas J (American Fisheries Society, Bethesda, MD), Vol 41, pp 325–343.
- Collie JS, Hall SJ, Kaiser MJ, Poiner IR (2000) A quantitative analysis of fishing impacts on shelf-sea benthos. *J Anim Ecol* 69:785–798.
- Kaiser MJ, et al. (2006) Global analysis and prediction of the response of benthic biota to fishing. *Mar Ecol Prog Ser* 311:1–14.
- van Denderen PD, et al. (2015) Similar effects of bottom trawling and natural disturbance on composition and function of benthic communities across habitats. *Mar Ecol Prog Ser* 541:31–43.
- Ramsay K, Kaiser MJ, Moore PG, Hughes RN (1997) Consumption of fisheries discards by benthic scavengers: Utilization of energy subsidies in different marine habitats. *J Anim Ecol* 66:884–896.
- Duplisea DE, Jennings S, Warr KJ, Dinmore TA (2002) A size-based model of the impacts of bottom trawling on benthic community structure. *Can J Fish Aquat Sci* 59: 1785–1795.
- Hiddink JG, et al. (2016) Bottom trawling affects fish condition through changes in the ratio of prey availability to density of competitors. *J Appl Ecol* 53:1500–1510.
- Rijnsdorp AD, Buys AM, Storbeck F, Visser EG (1998) Micro-scale distribution of beam trawl effort in the southern North Sea between 1993 and 1996 in relation to the trawling frequency of the sea bed and the impact on benthic organisms. *ICES J Mar Sci* 55:403–419.
- Lambert GI, et al. (2014) Quantifying recovery rates and resilience of seabed habitats impacted by bottom fishing. *J Appl Ecol* 51:1326–1336.
- Ellis N, Pantus F, Pitcher CR (2014) Scaling up experimental trawl impact results to fishery management scales—a modelling approach for a "hot time." *Can J Fish Aquat Sci* 71:733–746.
- Hiddink JG, Jennings S, Kaiser MJ (2006) Indicators of the ecological impact of bottom-trawl disturbance on seabed communities. *Ecosystems* 9:1190–1199.
- Schaefer MB (1954) Some aspects of the dynamics of populations important to the management of the commercial marine fisheries. *Inter-American Tropical Tuna Commission Bull* 1:23–56.
- McClanahan TR, Graham NA, Calnan JM, MacNeil MA (2007) Toward pristine biomass: Reef fish recovery in coral reef marine protected areas in Kenya. *Ecol Appl* 17: 1055–1067.
- Costello C, et al. (2016) Global fishery prospects under contrasting management regimes. *Proc Natl Acad Sci USA* 113:5125–5129.
- Pitcher CR, et al. (2017) Estimating the sustainability of towed fishing-gear impacts on seabed habitats: A simple quantitative risk assessment method applicable to data-poor fisheries. *Methods Ecol Evol* 8:472–480.
- Hiddink JG, et al. (2006) Cumulative impacts of seabed trawl disturbance on benthic biomass, production and species richness in different habitats. *Can J Fish Aquat Sci* 63: 721–736.
- Hughes KM, et al. (2014) Investigating the effects of mobile bottom fishing on benthic biota: A systematic review protocol. *Environ Evidence* 3:23.
- Baird S, Wood BA, Bagley NW (2011) *Nature and Extent of Commercial Fishing Effort on or Near the Seafloor Within the New Zealand 200 N. Mile Exclusive Economic Zone, 1989–90 to 2004–05* (Ministry of Fisheries, Wellington, New Zealand).
- Pitcher R, et al. (2016) *Implications of Current Spatial Management Measures for AFMA ERAs for Habitats*. (CSIRO Oceans & Atmosphere, Brisbane, Australia), FRDC Project 2014/204.
- Bolam SG, et al. (2017) Differences in biological traits composition of benthic assemblages between unimpacted habitats. *Mar Environ Res* 126:1–13.
- Tillin HM, Hiddink JG, Kaiser MJ, Jennings S (2006) Chronic bottom trawling alters the functional composition of benthic invertebrate communities on a sea basin scale. *Mar Ecol Prog Ser* 318:31–45.
- Brown CJ, Trebilco R (2014) Unintended cultivation, shifting baselines, and conflict between objectives for fisheries and conservation. *Conserv Biol* 28:677–688.
- Jennings S, Greenstreet SPR, Reynolds JD (1999) Structural change in an exploited fish community: A consequence of differential fishing effects on species with contrasting life histories. *J Anim Ecol* 68:617–627.
- Eigaard OR, et al. (2015) Estimating seabed pressure from demersal trawls, seines, and dredges based on gear design and dimensions. *ICES J Mar Sci* 73:i27–i43.
- Zuur AF, et al. (2009) *Mixed Effects Models and Extensions in Ecology with R* (Springer, New York).
- Pinheiro J, Bates D, DebRoy S, Sarkar D (2013) nlme: Linear and Nonlinear Mixed Effects Models. (R Package), Version 3.1-113.
- Jennings S, Lee J, Hiddink JG (2012) Assessing fishery footprints and the tradeoffs between landings value, habitat sensitivity and fishing impacts to inform marine spatial planning and an ecosystem approach. *ICES J Mar Sci* 69:1053–1063.
- Behrenfeld MJ, Falkowski PG (1997) Photosynthetic rates derived from satellite-based chlorophyll concentration. *Limnol Oceanogr* 42:1–20.
- Lutz MJ, Caldeira K, Dunbar RB, Behrenfeld MJ (2007) Seasonal rhythms of net primary production and particulate organic carbon flux to depth describe the efficiency of biological pump in the global ocean. *J Geophys Res Oceans* 112:C10011.
- Folk RL (1954) The distinction between grain size and mineral composition in sedimentary-rock nomenclature. *J Geol* 62:344–359.

Supporting Information

Text S1. Elaboration of systematic review process

Data were collated from published experimental and comparative studies of the effects of bottom trawling on seabed habitat and biota following a systematic review protocol (1).

‘Bottom trawling’ is defined here to include any commercial towed bottom gear, including otter trawls, beam trawls, scallop dredges and hydraulic dredges used to catch fish and invertebrates living in, on or in close association with seabed habitats. For the systematic review we attempted to find every study in journal papers, book chapters and grey literature reports that reported the effect of bottom trawling on the state of seabed (benthic) taxa (species or higher taxonomic levels) and communities (biomass, abundance, taxon richness and diversity). Each identified study had to pass quality assurance criteria before associated data were included in our analysis (1). This approach eliminated the possibility of bias in selection of the studies. We included studies that reported the effect of bottom fishing activities (exposure) on marine benthic biota (subject) and compared this with effects of no exposure or less exposure to bottom fishing gear (comparator). Studies also had to report a measurable effect (or outcome, non–significant results were included) on at least one identified component of the benthic biota (and to report outcomes from two or more areas of the seabed subject to different intensities of fishing disturbance. Data on the state of benthic biota were extracted from figures, tables or text within publications. We only used studies reporting whole community biomass and/or abundance of macrofaunal invertebrates (infauna and/or epifauna). This includes all species that were effectively sampled, including scavenging species. If essential data were missing, incomplete or contained obvious errors, the lead author was contacted to request these data and these data were included in the analysis if received. 42% of contacted authors responded and provided the requested data, 13% responded but could not supply the requested data, and 44% of authors did not respond. Meta–data were also extracted for each study (including location, depth, gear type, habitat, Table S2 & S3).

Most existing knowledge about fishing impacts has been gained from experimental studies, where abundance of benthic biota is recorded before and after experimental trawling. These studies were used to quantify the direct depletion d (or mortality) caused by the pass of a trawl (SI Text S2 for details on the analysis of this data). Comparative (or ‘gradient’ or ‘observational’) studies are studies where the benthic community is compared over a gradient of two or more levels of quantified fishing effort, where trawling effort may have been continuous, seasonal or a single event (SI Text S3

for details on the analysis of these data). The comparative studies allow the estimation of the ratio of d to r , and estimation of r when d is known from the experimental studies.

Gear types in the studies were classified as otter trawls (OT), beam trawls (BT), towed dredges (TD) and hydraulic dredges (HD). Otter trawls are widely used in all types of fisheries on a wide range of sediments and target species like gadoids, some flatfishes and prawns. The use of beam trawls is more restricted to sandy and gravelly bottoms and these gears are mostly used to target flatfishes and shrimps. Towed dredges are generally used to target scallops or other bivalve molluscs, and are often fished on gravelly bottoms. Hydraulic dredges are used to target buried bivalves and resuspend sediment to a depth of up to 40 cm. They are often used in intertidal and other shallow areas (2, 3).

Text S2. Estimating depletion from experimental studies and penetration depth

Depletion d for macrofauna community biomass and abundance was estimated from the experimental studies identified in the systematic review. Studies used before–after (BA), control–impact (CI), or before–after–control–impact (BACI) designs. 13 of the studies were carried out in areas that were trawled in the last two years but generally at low intensity, 9 were carried out in areas that were last trawled between 10 and 25 years ago, and 10 were carried out in areas that have no fishing history. For the remaining studies this information was not given. Most experiments have therefore been conducted in infrequently trawled and untrawled areas, this is possible because even in the most heavily trawled areas like Europe about one-third of the seabed is not trawled (3). We used the log response ratio ($\ln RR$) as the response variable, which was calculated as $\ln(\text{response fished} / \text{response control})$ for CI studies and $\ln(\text{response after} / \text{response before})$ for BA studies. The combined variance was calculated as in Borenstein et al. (4). For BACI studies, calculation of the $\ln RR$ and combined variance was more complicated. Let $y = \log X$ denote the log abundance and consider the four combinations of control/impact and before/after: y_{CB} , y_{CA} , y_{IB} , and y_{IA} . We assumed that effects are multiplicative on the abundance scale, and therefore are additive on the log scale. Let a be the before–after (period) effect, b the treatment effect, and c the interaction term. Then $y_{CB} = \mu$, $y_{CA} = \mu + a$, $y_{IB} = \mu + b$ and $y_{IA} = \mu + a + b + c$.

This means that $c = y_{IA} - y_{IB} - y_{CA} + y_{CB}$. On the abundance scale this implies

$$c = \log \left[\frac{X_{IA}}{X_{IB}} / \frac{X_{CA}}{X_{CB}} \right] \quad (\text{eq S2.1})$$

The quantity c is the analogue of $\ln RR$ for BACI data. The variance calculation uses the following approximation:

$$\text{Var}[\log X] \simeq \frac{\text{Var}[X]}{[EX]^2} \quad (\text{eq S2.2})$$

This leads to this expression for the variance

$$\text{Var}[c] \simeq \frac{SD_{IA}^2}{n_{IA}\bar{X}_{IA}^2} + \frac{SD_{IB}^2}{n_{IB}\bar{X}_{IB}^2} + \frac{SD_{CA}^2}{n_{CA}\bar{X}_{CA}^2} + \frac{SD_{CB}^2}{n_{CB}\bar{X}_{CB}^2} \quad (\text{eq S2.3})$$

The $\ln RR$ will be more negative in areas that have been exposed to a higher frequency of fishing disturbance. Therefore, it was corrected using $\ln RR = \ln RR_{\text{uncorrected}} / Idis$ where $Idis$ is the number of trawl passes over the fished area.

The number of data points available for estimating d was limited: 55 for community biomass and 101 for community abundance (OT: 25, BT: 6, TD: 87, HD: 38). Including the response unit (biomass or

abundance) as a factor in the model did not result in a lower AIC, therefore we estimated d using both biomass and abundance values in a single model.

Penetration depth of different gear types

Predicted penetration depth (P) of each trawl type was estimated from the penetration depth of the individual components of the gear weighted by the width of these components. Penetration depth is defined as the depth to which the sediment was disturbed by the fishing gear, but in practice often measured as the depth to which the sediment was excavated. We conducted a systematic search of the literature starting from Table 6 in Eigaard et al. (5). Each reference in the table was checked and only included when a study directly measured penetration depth. A database of experimental and comparative studies of fishing impacts, produced during a systematic review (1) was also screened for further studies that provided measurements of penetration depth. In addition, references cited within each reference already identified were screened and further studies included as a result. Any study for which penetration depth of a fishing gear (whole), or a gear component, was measured or inferred by one of the following methods was included: underwater video, underwater photographs side-scan sonar, sediment profile images, markers in sediment, observations by SCUBA divers, high resolution acoustic array, underwater laser, inferred from the living position of benthic organisms retained by the fishing gear, or in the case of intertidal fishing methods – by direct observation. Because different methods were used, estimates of penetration depths across studies may not be directly comparable, although they are the best available estimates. Review papers that were not the primary source of penetration depth data were not included, but were used to identify primary sources of data. Studies that reported penetration depths but that were not included in our analysis are given in Table S8.

The sources we identified reported the penetration depth either for the whole gear or for individual gear components (e.g., doors, sweeps, and bridles of an OT). The predicted penetration depth per gear component was therefore estimated by fitting a nested linear model where $\log(\text{penetration}+1) \sim \text{sediment type} + \text{Gear} | \text{Component}$. Although we were not directly interested in the effect of sediment type, it was included because within gears the penetration seemed to vary with sediment type and it allowed us to correct for this effect in the final P estimates (Gear:Component $F_{9,74}=6.57$, $p < 0.001$, Habitat $F_{3,71}=2.6$, $p=0.057$). We used the fitted model to predict the penetration depth for each gear component in each sediment type, and estimated the overall P for each fishing gear from

107 this in two steps. We first averaged predicted penetration depths over all sediment types for each
108 gear component, and then estimated the mean P for each gear by taking the mean weighted by the
109 width of these components.

Text S3. Estimating the effect of trawling in comparative studies

As described in the main text, the responses from different studies were normalised to the common units of B/K . K was estimated for each of the sampling methods as $10^{\text{intercept}}$ of the relationship of $\log_{10} B$ versus F . In some studies the biota was sampled using two or more methods, each suited to sampling a different component of the community. For example, Hiddink et al. (6) sampled each station across a gradient of trawling frequency with an anchor dredge, box corer and 2m beam trawl. Where two or more sampling methods were used to sample benthic community biomass, K was estimated separately for each sampling gear. Studies were treated as replicate measurements by using study as a random effect.

A collective analysis of gradient studies requires fishing pressure to be described on a common scale. We adopted trawling frequency, F (y^{-1}), which is equivalent to the swept area ratio ($\text{km}^2 \text{ km}^{-2} \text{ y}^{-1}$). Trawling frequency expresses how often each cell is trawled in a year, and is calculated by dividing the area trawled in a year by the area of the study site or other defined area (e.g. grid cell). Trawled area is usually calculated using logbook or vessel monitoring system (VMS) data, from the number of hours spent fishing multiplied by the fishing speed and the width of the fishing gear. Trawling frequency was explicitly reported for about half the comparative studies, and for the other half we calculated trawling frequency from the reported fishing effort (Table S9). Where trawling frequency could not be calculated, the study was excluded from further analyses.

Here we apply eq. 3.1 for estimating the effect of trawling on B/K for groups of species and communities. These communities, however, comprise many species with wide variety of r and K values. Therefore, the response to fishing is the sum of the responses of all those species. Because low- r species will be more depleted than high- r species, and will potentially be extirpated from the community, the response of the community to F is not a straight line as in eq. 3.1. Consequently, the average r of the community increases with F , and the marginal effect of each additional unit of F on community B/K decreases with increasing F . We simulated a community of species by drawing r and K values at random, and found that the resulting relationship between total community B and F is well approximated by a log-linear relationship for normal and exponential distributions of r and K . We therefore estimated the effect of trawling on communities by fitting a model based on the approximation:

$$\log_{10}(B/K) \sim bF \quad (\text{eq. S3.1})$$

143 where b is the slope of the relationship. After fitting a linear relationship to $\log_{10} B$ versus F for each
144 comparative study, K was estimated as the $10^{\text{intercept}}$ of this relationship.

Text S4. Estimating r from d and b and quantifying uncertainty

Comparative studies involve sampling the seabed biota at locations within sites subject to different frequencies of trawling disturbance. Collectively, the sampling locations only cover a small proportion of each site, but the mean trawling frequency estimated for the site is assumed to apply to all stations within the site because data on trawl positions are not sufficiently resolved to estimate location-specific trawling frequency. Thus samples linked to the same mean trawling frequency for the site may come from heavily trawled patches, lightly trawled patches, and potentially some untrawled patches, within the site. Consequently, the mean recovery rate estimated for the site (R) will not be the same as the intrinsic rate of recovery r in equation (1). If the distribution which describes the patchiness of trawling within a site is known then r can be estimated following the approach in Ellis et al. (7). Given that $\log_{10}(B/K) \sim bF$ (SI Text S3) and that $B/K = 1 - (d/R) F$, it follows that $10^{bF} = 1 - (d/R) F$. The equation to estimate R from r for a single species in Ellis et al. (7, $R = r \log(1+\beta d)/[-\beta \log(1-d)]$) can therefore be rewritten to estimate r for the community as:

$$r = R / \frac{\log(1+\beta d)}{-\beta \log(1-d)} \quad (\text{eq S4.1})$$

where

$$R = \frac{-d}{(10^{bF} - 1)/F} \quad (\text{eq S4.2})$$

where β is a parameter defining the spatial distribution of trawling within a site (7). Here we assumed $\beta \approx 0$, representing a random distribution of trawling within a site (in practice $\beta = 10^{-6}$ because the equation is undefined when $\beta = 0$). A random distribution within sites is supported by data on the spatial distribution of trawling collected at scales of around 1 km and smaller (8), consistent with the scales at which sites in comparative trawling studies are defined. Assuming a uniform distribution of trawling ($\beta = -1$) resulted in r estimates that were approximately 10% lower. Equation S4.1 and S4.2 indicate that r depends on F , which is expected because changes in community composition to favour biota with faster life histories. Because we aim to estimate recovery rates and times for the original unfished community, we used estimates of r at $F = 0$ to estimate recovery times. If the distribution of trawling in a cell is random, the site level depletion is the same as local depletion d and no correction was therefore applied here.

176 To propagate the uncertainty in the estimates of b and d into the estimate of r we sampled the
177 distributions of b and d estimates to derive the distribution of r . The value of b was taken as negative
178 and $-b$ was assumed to have a log-normal distribution, with the standard deviation estimated from
179 the distribution of the random slopes using the *fitdist* function in the *fitdistrplus* package in R (9).
180 The value of d was assumed to be positive and bounded between 0 and 1, and to have a logitnormal
181 distribution with standard deviation estimated with the function *twCoefLogitnorm* of the *logitnorm*
182 package in R (10). We sampled 2000 combinations from the distributions of b and d to estimate the
183 distribution of r .

Text S5. Estimating recovery time from r

The logistic r can be used to estimate recovery time (i.e. T from a defined level of depletion below K to a defined proportion of K). Lambert et al. (11) derived the recovery time T to $0.9K$ as:

$$T = \frac{1}{r} \left[\ln \left(\frac{0.9K}{B_{t=0}} \right) + \ln \left(\frac{K - B_{t=0}}{0.1K} \right) \right] \quad (\text{eq. S5.1})$$

If we generalise this in terms of any fraction of K at which recovery is deemed to have occurred (ϕ) and assume that $B_{t=0}$ is the biomass or abundance of an unimpacted habitat remaining after the pass of a gear that reduces biomass or abundance by a fraction d , then the recovery time given these conditions would be:

$$T = \frac{1}{r} \left[\ln \left(\frac{\phi K}{K(1-d)} \right) + \ln \left(\frac{K - K(1-d)}{K(1-\phi)} \right) \right] \quad (\text{eq. S5.2})$$

which can be expressed more simply as:

$$T = \frac{1}{r} \ln \left(\frac{\phi d}{(1-d)(1-\phi)} \right) \quad (\text{eq. S5.3})$$

Table S1. Number of studies of whole community biomass and abundance for macrofauna per gear and habitat. Otter trawls (OT), beam trawls (BT), towed dredges (TD), hydraulic dredges (HD).

a) Experimental studies

	OT	BT	TD	HD
Biogenic	–	–	–	–
Gravel	1	–	1	–
Sand	6	4	16	10
Sandy mud/Muddy sand	–	–	–	2
Mud	5	–	–	1

b) Comparative studies

	OT	BT	TD	HD
Biogenic	–	–	–	–
Gravel	1	–	5	–
Sand	3	4	–	–
Sandy mud/Muddy sand	5	2	–	–
Mud	4	–	–	–

211 Table S2. Metadata for included experimental studies. A single paper is listed more than once when
 212 two or more studies were reported in the same paper.

Source	Region	Habitat	Depth (m)	Gear
(12)	Southern Europe	S	8	TD
(12)	Southern Europe	S	8	TD
(13)	Alaska	S	25	OT
(14)	Southern Europe	S	9	TD
(14)	Southern Europe	S	9	TD
(15)	Southern Europe	S	6	TD
(15)	Southern Europe	S	18	TD
(16)	Australia	S	20	OT
(16)	Australia	M	18	OT
(16)	Australia	S	20	OT
(17)	Northern Europe	S	10	TD
(17)	Northern Europe	S	10	TD
(17)	Northern Europe	S	10	TD
(17)	Northern Europe	S	10	TD
(18)	North America	mS	65	HD
(19)	North America	S	5.5	HD
(20)	Northern Europe	mS	0	HD
(21)	Northern Europe	S	7	HD
(22)	North America	G	70	OT
(23)	Northern Europe	S	21.5	TD
(23)	Northern Europe	S	21.5	OT
(23)	Northern Europe	S	21.5	TD
(24)	Northern Europe	S	26	BT
(24)	Northern Europe	S	34	BT
(25)	Northern Europe	S	0	HD
(26)	Northern Europe	S	30	BT
(26)	Northern Europe	S	30	BT
(27)	North America	S	0.2	HD
(27)	North America	S	0.2	HD
(27)	North America	S	0.2	HD
(28)	Southern Europe	S	24	TD

(29)	Southern Europe	S	23	TD
(29)	Southern Europe	M	11	TD
(30)	South America	S	10	OT
(31)	Canada	S	133	OT
(32)	Southern Europe	M	30	OT
(32)	Southern Europe	M	40	OT
(33)	Australia	S	0	HD
(34)	North America	M	61	OT
(35)	Northern Europe	M	0	HD
(36)	New Zealand	S	24	TD
(37)	New Zealand	S	24	TD
(38)	Northern Europe	M	33.5	OT
(39)	Northern Europe	S	3.5	HD
(40)	South Africa	S	0	HD
(40)	South Africa	S	0	HD

213

214

215 Table S3. Metadata for included comparative studies. sM & mS – sandy mud and muddy sand.

Source	Region	Habitat	Depth (m)	Gear
(41)	South Africa	sM & mS	420	OT
(42)	Eastern North America	Gravel	48	TD
(43)	South Coast Australia	Sand	30	OT
(44)	North Sea	Mud	80	OT
(6)	North Sea	Sand	32.5	BT
(6)	North Sea	Sand	40	BT
(45)	Northwest Europe	sM & mS	31.5	OT
(46)	North west Europe	sM & mS	31.5	OT
(47)	Central North Sea,	sM & mS	57.5	BT
(47)	Central North Sea,	Sand	57.5	BT
(48)	North Sea	sM & mS	50	BT
(49)	Irish Sea	Gravel	43.5	TD
(50)	Mediterranean Sea	Mud	137.145	OT
(51, 52)*	Australia	sM & mS	27.5	OT
(51, 52)*	Australia	Sand	25.5	OT
(53)	Irish Sea	sM & mS	30	OT
(54)	North West Europe	Sand	40	BT
(55)	Eastern North America	Gravel	74	TD
(55)	Eastern North America	Gravel	50	TD
(56)	Australia	Sand	23.5	OT
(57)	North west Europe	Mud	147.5	OT
(57)	North west Europe	Gravel	78.5	OT
(58)	Irish Sea	Gravel	43.5	TD
(59)	North west Europe	Mud	100	OT

216 * sources combined

217

218 Table S4. Penetration depth P and depletion d of community biomass and abundance for different
 219 trawling gears. The 5 and 95% percentiles for d estimates are given. Gear types are otter trawls (OT),
 220 beam trawls (BT), towed dredges (TD) and hydraulic dredges (HD).

Gear	Penetration depth (cm)			Depletion d (fraction)		
	mean	±	sd	5%	Median	95%
OT	2.44	±	1.14	0.02	0.06	0.16
BT	2.72	±	1.24	0.07	0.14	0.25
TD	5.47	±	2.19	0.13	0.20	0.30
HD	16.11	±	5.80	0.35	0.41	0.48

221

222

223

224 Table S5. AIC estimates of the linear mixed models with different explanatory variables for
 225 community biomass and abundance in comparative studies. The model with the lowest AIC for
 226 biomass and the two models with the lowest AIC for abundance are given in bold.

Model	Biomass	Abundance
None	566.9	89.5
Habitat	573.1	94.3
Gear	572.5	92.4
<i>d</i>	568.9	89.5
Penetration	568.8	89.3
SBT	568.8	89.1
Depth	567.7	91.4
POC	567.7	91.1
PP	568.0	90.9
Gravel	568.4	81.1
Sand	568.8	89.2
Mud	568.9	90.2
<i>d</i> /SBT	568.9	89.5
<i>d</i> *Depth	567.9	90.8
<i>d</i> /POC	568.7	91.1
<i>d</i> /PP	568.8	86.1
Penetration/SBT	568.9	90.0
Penetration×Depth	567.8	91.4
Penetration/POC	568.5	91.4
Penetration/PP	568.7	89.0

227

228 d = depletion estimate from experimental studies (fraction per trawl pass)

229 Penetration = penetration depth of fishing gear into the seabed (cm)

230 SBT = sea bottom temperature ($^{\circ}\text{C}$)

231 POC = Particulate organic carbon flux to the seabed ($\text{g C}_{\text{org}} \text{ m}^{-2} \text{ yr}^{-1}$)

232 PP = Primary production ($\text{mg C m}^{-2} \text{ d}^{-1}$)

233 Gravel, Sand & Mud = sediment composition in % by weight

234 Habitat = categorical variable with levels Mud, sM & mS, Sand and Gravel.

235

236

237 Table S6. Parameters used to estimate r and percentiles from the distribution of r estimates. SD =
 238 standard deviation. SAR= swept area ratio.

	Biomass	Abundance		
Gear type	Combined	OT	BT	TD
Gravel content (%)	NA	0.84	0.00	44.63
d (fraction)	0.13	0.06	0.14	0.20
SD of d	0.08	0.08	0.08	0.08
b (slope)	-0.075	-0.025	-0.015	-0.553
SD of b	0.003	0.057	0.057	0.057
% decline with unit increase in SAR	15.90	5.50	3.29	71.99
Recovery time from 0.5K to 0.95K (years, using median r)	3.58	2.81	0.66	16.65
r , 5% percentile	0.42	0.33	2.37	0.11
r , 10% percentile	0.49	0.43	2.75	0.12
r , 25% percentile	0.63	0.66	3.50	0.15
r , 50% percentile	0.82	1.05	4.49	0.18
r , 75% percentile	1.06	1.66	5.71	0.21
r , 90% percentile	1.34	2.60	7.18	0.25
r , 95% percentile	1.54	3.54	8.21	0.28

239

240

241 Table S7. Studies used to estimate the penetration depth of different gear types. Weighting is the
 242 fraction of the width of the gear occupied by a component.

Source	Gear	Component	Habitat	Penetration (cm)	Weighting
(60)	BT	Beam trawl - whole gear	Sand	6	1
(61)	BT	Beam trawl - whole gear	Sand	2.26	1
(61)	BT	Beam trawl - whole gear	Mud	5.29	1
(62)	BT	Beam trawl - whole gear	Sand	1	1
(63)	BT	Beam trawl - tickler chains	Mud	1.4	0.94
(64)	BT	Beam trawl - tickler chains	Sand	0.75	0.94
(65)	BT	Beam trawl - tickler chains	Sand	6	0.94
(63)	BT	Beam trawl - tickler chains	Sand	0.4	0.94
(63)	BT	Beam trawl - tickler chains	Gravel	0.5	0.94
(64)	BT	Beam trawl - tickler chains	Mud	0.9	0.94
(61)	BT	Beam trawl - tickler chains	Sand	1	0.94
(64)	BT	Beam trawl - tickler chains	Sand	0	0.94
(61)	BT	Beam trawl - trawl shoes	Sand	1.9	0.06
(64)	BT	Beam trawl - trawl shoes	Sand	1.5	0.06
(66)	OT	Otter trawl - whole gear	Mud	8.5	1
(67)	OT	Otter trawl - whole gear	Sand	4.5	1
(66)	OT	Otter trawl - whole gear	Sand	0.085	1
(68)	OT	Otter trawl - whole gear	Gravel	4.5	1
(69)	OT	Otter trawl - sweeps	Mud	2.18	0.73
(69)	OT	Otter trawl - ground gear	Mud	1.4	0.25
(70)	OT	Otter trawl - ground gear	Mud	0	0.25
(71)	OT	Otter trawl - trawl doors	Mud	30	0.02
(72)	OT	Otter trawl - trawl doors	Mud	12.5	0.02
(73)	OT	Otter trawl - trawl doors	Mud	5.5	0.02
(73)	OT	Otter trawl - trawl doors	Sand	2.7	0.02
(69)	OT	Otter trawl - trawl doors	Mud	6.43	0.02
(71)	OT	Otter trawl - trawl doors	Sand	20	0.02
(72)	OT	Otter trawl - trawl doors	Sand	2.5	0.02
(74)	OT	Otter trawl - trawl doors	Sand	10	0.02
(75)	OT	Otter trawl - trawl doors	Mud	5	0.02
(75)	OT	Otter trawl - trawl doors	Sand	2.5	0.02
(69)	OT	Otter trawl - trawl doors	Sand	0.26	0.02
(69)	OT	Otter trawl - trawl doors	Sand	2.1	0.02

(69)	OT	Otter trawl - trawl doors	Sand	5.8	0.02
(69)	OT	Otter trawl - trawl doors	Sand	0.2	0.02
(76)	OT	Otter trawl - trawl doors	Gravel	5.5	0.02
(77)	OT	Otter trawl - trawl doors	Sand	15	0.02
(62)	OT	Otter trawl - trawl doors	Mud	14	0.02
(70)	OT	Otter trawl - trawl doors	Mud	4.5	0.02
(75)	OT	Twin Otter trawl - roller clump	Sand	0	0.01
(75)	OT	Twin Otter trawl - roller clump	Mud	0	0.01
(72)	OT	Twin Otter trawl - roller clump	Mud	3.5	0.01
(73)	OT	Twin Otter trawl - roller clump	Mud	12.5	0.01
(76)	OT	Twin Otter trawl - roller clump	Mud	12	0.01
(73)	OT	Twin Otter trawl - roller clump	Sand	3.65	0.01
(66)	OT	Otter trawl - whole gear	Mud	8.5	1
(67)	OT	Otter trawl - whole gear	Sand	4.5	1
(66)	OT	Otter trawl - whole gear	Sand	0.085	1
(68)	OT	Otter trawl - whole gear	Gravel	4.5	1
(69)	OT	Otter trawl - sweeps	Mud	2.18	0.73
(69)	OT	Otter trawl - ground gear	Mud	1.4	0.25
(70)	OT	Otter trawl - ground gear	Mud	0	0.25
(71)	OT	Otter trawl - trawl doors	Mud	30	0.01
(72)	OT	Otter trawl - trawl doors	Mud	12.5	0.01
(73)	OT	Otter trawl - trawl doors	Mud	5.5	0.01
(73)	OT	Otter trawl - trawl doors	Sand	2.7	0.01
(69)	OT	Otter trawl - trawl doors	Mud	6.43	0.01
(71)	OT	Otter trawl - trawl doors	Sand	20	0.01
(72)	OT	Otter trawl - trawl doors	Sand	2.5	0.01
(74)	OT	Otter trawl - trawl doors	Sand	10	0.01
(75)	OT	Otter trawl - trawl doors	Mud	5	0.01
(75)	OT	Otter trawl - trawl doors	Sand	2.5	0.01
(69)	OT	Otter trawl - trawl doors	Sand	0.26	0.01
(69)	OT	Otter trawl - trawl doors	Sand	2.1	0.01
(69)	OT	Otter trawl - trawl doors	Sand	5.8	0.01
(69)	OT	Otter trawl - trawl doors	Sand	0.2	0.01
(76)	OT	Otter trawl - trawl doors	Gravel	5.5	0.01
(77)	OT	Otter trawl - trawl doors	Sand	15	0.01
(62)	OT	Otter trawl - trawl doors	Mud	14	0.01
(70)	OT	Otter trawl - trawl doors	Mud	4.5	0.01
(35)	HD	Hydraulic dredge	Mud	10	1

(18)	HD	Hydraulic dredge	Sand	20	1
(19)	HD	Hydraulic dredge	Sand	5	1
(20)	HD	Hydraulic dredge	Mud	30	1
(21)	HD	Hydraulic dredge	Sand	25	1
(25)	HD	Hydraulic dredge	Mud	10	1
(78)	HD	Hydraulic dredge	Sand	9	1
(79)	HD	Hydraulic dredge	Sand	40	1
(20)	HD	Tractor dredge	Mud	30	1
(80)	HD	Hydraulic dredge	Sand	5	1
(28)	TD	Boat Dredge - whole gear	Sand	6	1
(17)	TD	Boat Dredge - teeth	Sand	3.5	1
(81)	TD	Boat Dredge - teeth	Maerl	10	1
(82)	TD	Boat Dredge - whole gear	Sand	2.5	1
(82)	TD	Boat Dredge - whole gear	Gravel	3.5	1
(82)	TD	Boat Dredge - whole gear	Gravel	5.9	1

243

244

Table S8. A list of studies which relate to the physical impacts of towed-bottom fishing gears, but were not included in the penetration depth calculations for the reasons given.

Source	Reason for non-inclusion
(83)	Not the primary source of the reported gear penetration depth
(84)	Penetration depth not reported
(85)	Penetration depth not reported
(86)	Penetration depth not reported
(87)	Review paper, therefore not a primary data source
(88)	Penetration depths obtained from a numerical model, rather than direct measurements
(89)	Penetration depth not reported
(90)	Not the primary source of the reported gear penetration depth
(91)	Not the primary source of the reported gear penetration depth
(92)	Unable to obtain manuscript, however a penetration depth of c. 6.5cm is cited in de Groot (1995) and referenced to this report.
(93)	Not the primary source of the reported gear penetration depth
(94)	The source of the seabed marks measured in the study is ambiguous
(95, 96)	Penetration depth inferred from amount of suspended sediment only, missing non-suspended component of penetration.

Table S9. Trawling frequency calculations for comparative studies where trawling frequency was not reported as the swept area ratio (SAR).

Paper	Region	Habitat	Depth (m)	Gear	Gear	Source of swept area ratio estimate (SA)	Reported effort	Area box	Fishing speed	Gear width (m)	SAR calculation (after converting to the same units)	Min SAR (y^{-1})	Max SAR (y^{-1})
Abbreviation							E	A	Sp	W			
Collie et al. 2005	Georges Bank, North America	Gravel	48	Scallop dredge	TD	Calculate d	hrs fished y^{-1}	1 nm ²	3 kn	8	E*Sp*W/A	0.0	3.7
Currie et al. 2011	Spencer Gulf , South Eastern Australia	Sand	30	Prawn trawl	OT	Calculate d	h km ⁻²		3 kn	29.26	E*Sp*W	0.1	2.9
Frid et al. 1999	Northumberland, NE England, USA	Mud	80	Otter trawls	OT	Calculate d	km ² trawled y^{-1}	ICES++			E/A	0.0	12.9
Jennings et al. 2001a	Silver Pit, North Sea	Sm & Ms	57.5	Beam trawl	BT	Estimate d using VMS						0.5	5.4
Jennings et al. 2001b	Hills, North Sea	Sand	57.5	Beam trawl	BT	Estimate d using VMS						0.1	2.3
Jennings et al. 2002	Silver Pit, North Sea	Sm & Ms	50	Beam and otter trawls	BT	Estimate d using VMS						0.4	5.0
Kaiser et al.	Irish Sea, Isle of	Gravel	43.5	Scallop	TD	Calculate	hrs fished	5x5 nm	2.5 kn	12.0†	E*Sp*W/A	0.1	3.2

2000b	Man			dredge		d	y ⁻¹							
Reiss et al.	German Bight,	Sand	40	Beam	BT	Reported							0.1	2.0
2009	Germany			trawl										
Smith et al.	North East	Gravel	74	Scallop	TD	Calculate	hrs fished	50 km ²	3 kn	30	E*Sp*W/A	0.0	0.4	
2013a	Peak, Georges			dredge		d	y ⁻¹							
	Bank, N.													
	America													
Smith et al.	CAI, Georges	Gravel	50	Scallop	TD	Calculate	hrs fished	50 km ²	3 kn	30	E*Sp*W/A	0.0	0.4	
2013b	Bank, N.			dredge		d	y ⁻¹							
	America													
Svane et al.	Spencer Gulf,	Sand	23.5	Prawn	OT	Calculate	h trawled	645 to	2.5 kn	29.26	E*Sp*W/A	0.2	1.9	
2009	Australia			trawl		d	per year	1128						
								km2						
Veale et al.	Irish Sea, Isle of	Gravel	43.5	Scallop	TD	Calculate	m x h y ⁻¹	5x5 nm	2.5 kn	10	E*Sp*W/A	0.0	1.7	
2000	Man			dredge		d								
Vergnon &	Grande Vasiere,	Mud	100	Nephrops	OT	Calculate	mths	ICES	2 kn	10	E*Sp*W/A	1.6	7.9	
Blanchard	Bay of Biscay,			trawl		d	fished y ⁻¹	rectangl						
2006	France							e (1 *						
								0.5						
								degree)						

† 16 dredges (10 within 3nmi limit) of 0.75m width.

†† ICES rectangles of size 1° × 0.5°.




Figure S1. Predicted increase in median r (± 5 -95% quantiles) with trawling frequency for community biomass, as estimated from the relationship between $\log_{10}B/K$ and trawling frequency (equations S4.1 and S4.2). Simulation assumes mean value of d over all fishing gears included in the comparative studies ($d = 0.13$).

References

1. Hughes KM, et al. (2014) Investigating the effects of mobile bottom fishing on benthic biota: a systematic review protocol. *Environmental Evidence* 3:23:DOI: 10.1186/2047-2382-1183-1123
2. Sainsbury JC (1986) Commercial fishing methods. Fishing News Books, Oxford
3. Eigaard OR, et al. (2016) The footprint of bottom trawling in European waters: distribution, intensity and seabed integrity. *ICES J Mar Sci* 10.1093/icesjms/fsw194
4. Borenstein M, Hedges LV, Higgins JPT, Rothstein HR (2009) Introduction to Meta-Analysis. Wiley, Chichester, UK
5. Eigaard OR, et al. (2015) Estimating seabed pressure from demersal trawls, seines, and dredges based on gear design and dimensions. *ICES J Mar Sci* 73 S1:i27-i43.
6. Hiddink JG, et al. (2006) Cumulative impacts of seabed trawl disturbance on benthic biomass, production and species richness in different habitats. *Can J Fish Aquat Sci* 63:721-736
7. Ellis N, Pantus F, Pitcher CR (2014) Scaling up experimental trawl impact results to fishery management scales—a modelling approach for a “hot time”. *Can J Fish Aquat Sci* 71:733-746
8. Rijnsdorp AD, Buys AM, Storbeck F, Visser EG (1998) Micro-scale distribution of beam trawl effort in the southern North Sea between 1993 and 1996 in relation to the trawling frequency of the sea bed and the impact on benthic organisms. *ICES J Mar Sci* 55:403-419
9. Delignette-Muller ML, Dutang C (2015) fitdistrplus: An R Package for Fitting Distributions. *J Stat Softw* 64:1-34
10. Wutzler T (2012) logitnorm: Functions for the logitnormal distribution. R package version 0.8.29. <http://CRAN.R-project.org/package=logitnorm>.
11. Lambert GI, et al. (2014) Quantifying recovery rates and resilience of seabed habitats impacted by bottom fishing. *J Appl Ecol* 51:1326–1336
12. Alves F, Chícharo L, Nogueira A, Regala J (2003) Changes in benthic community structure due to clam dredging on the Algarve coast and the importance of seasonal analysis. *J Mar Biol Assoc UK* 83:719-729
13. Brown E, Finney B, Hills S, Dommissie M (2005) Effects of commercial otter trawling on benthic communities in the southeastern Bering Sea. *American Fisheries Society Symposium* 41:439-491
14. Carvalho S, et al. (2011) Relationship between razor clam fishing intensity and potential changes in associated benthic communities. *J Shellfish Res* 30:309-323

15. Constantino R, et al. (2009) Clam dredging effects and subsequent recovery of benthic communities at different depth ranges. *Mar Environ Res* 67:89-99
16. Drabsch SL, Tanner JE, Connell SD (2001) Limited infaunal response to experimental trawling in previously untrawled areas. *ICES J Mar Sci* 58:1261-1271
17. Eleftheriou A, Robertson MR (1992) The effects of experimental scallop dredging on the fauna and physical environment of a shallow sandy community. *Neth J Sea Res* 30:289-299
18. Gilkinson KD, et al. (2005) Immediate impacts and recovery trajectories of macrofaunal communities following hydraulic clam dredging on Banquereau, eastern Canada. *ICES J Mar Sci* 62:925-947
19. Goldberg R, et al. (2012) Effects of hydraulic shellfish dredging on the ecology of a cultivated clam bed. *Aquaculture Environment Interactions* 3:11-21
20. Hall SJ, Harding MJ (1997) Physical disturbance and marine benthic communities: the effects of mechanical harvesting of cockles on non-target benthic infauna. *J Appl Ecol* 34:497-517
21. Hall SJ, Basford DJ, Robertson MR (1990) The Impact of Hydraulic Dredging for Razor Clams *Ensis* Sp. on an Infaunal Community. *Neth J Sea Res* 27:119-125
22. Henry L-A, et al. (2006) Impacts of otter trawling on colonial epifaunal assemblages on a cobble bottom ecosystem on Western Bank (northwest Atlantic). *Mar Ecol Prog Ser* 306:63-78
23. Hinz H, Murray LG, Malcolm FR, Kaiser MJ (2012) The environmental impacts of three different queen scallop (*Aequipecten opercularis*) fishing gears. *Mar Environ Res* 73:85-95
24. Kaiser MJ, Spencer BE (1996) The effects of beam-trawl disturbance on infaunal communities in different habitats. *J Anim Ecol* 65:348-358
25. Kaiser MJ, Edwards DB, Spencer BE (1996) Infaunal community changes as a result of commercial clam cultivation and harvesting. *Aquat Living Resour* 9:57-63
26. Kaiser MJ, et al. (1998) Changes in megafaunal benthic communities in different habitats after trawling disturbance. *ICES J Mar Sci* 55:353-361
27. Peterson CH, Summerson HC, Fegley SR (1987) Ecological consequences of mechanical harvesting of clams. *Fish Bull* 85:281-298
28. Pranovi F, et al. (2000) Rapido trawling in the northern Adriatic Sea: effects on benthic communities in an experimental area. *ICES J Mar Sci* 57:517-524
29. Pranovi F, et al. (2005) Trawl Fishing Disturbance and Medium-Term Microfaunal Recolonization Dynamics: A Functional Approach to the Comparison between Sand and Mud Habitats in the Adriatic Sea (Northern Mediterranean Sea). In: American Fisheries Society Symposium, Vol 41. American Fisheries Society, p 545

30. Prantoni AL, et al. (2013) An experimental evaluation of the short-term effects of trawling on infaunal assemblages of the coast off southern Brazil. *J Mar Biol Assoc UK* 93:495-502
31. Prena J, et al. (1999) Experimental otter trawling on a sandy bottom ecosystem of the Grand Banks of Newfoundland: analysis of trawl bycatch and effects on epifauna. *Mar Ecol Prog Ser* 181:107-124
32. Sanchez P, Demestre M, Ramon M, Kaiser MJ (2000) The impact of otter trawling on mud communities in the northwestern Mediterranean. *ICES J Mar Sci* 57:1352-1358
33. Skilleter GA, Zharikov Y, Cameron B, McPhee DP (2005) Effects of harvesting callinassid (ghost) shrimps on subtropical benthic communities. *J Exp Mar Biol Ecol* 320:133-158
34. Sparks-McConkey PJ, Watling L (2001) Effects on the ecological integrity of a soft-bottom habitat from a trawling disturbance. *Hydrobiologia* 456:73-85
35. Spencer BE, Kaiser MJ, Edwards DB (1998) Intertidal clam harvesting: benthic community change and recovery. *Aquac Research* 29:429-437
36. Thrush SF, Hewitt JE, Cummings VJ, Dayton PK (1995) The impact of habitat disturbance by scallop dredging on marine benthic communities: what can be predicted from the results of experiments? *Mar Ecol Prog Ser* 129:141-150
37. Pikitch EK, et al. (2004) Ecosystem-based fishery management. *Science* 305:346-347
38. Tuck ID, et al. (1998) Effects of physical trawling disturbance in a previously unfished sheltered Scottish sea loch. *Mar Ecol Prog Ser* 162:227-242
39. Tuck ID, et al. (2000) The impact of water jet dredging for razor clams, *Ensis* spp., in a shallow sandy subtidal environment. *J Sea Res* 43:65-81
40. Wynberg RP, Branch GM (1994) Disturbance associated with bait-collection for sandprawns (*Callinassa kraussi*) and mudprawns (*Upogebia africana*): long-term effects on the biota of intertidal sandflats. *J Mar Res* 52:523-558
41. Atkinson LJ, Field JG, Hutchings L (2011) Effects of demersal trawling along the west coast of southern Africa: Multivariate analysis of benthic assemblages. *Mar Ecol Prog Ser* 430:241-255
42. Collie JS, Hermesen JM, Valentine PC, Almeida FP (2005) Effects of fishing on gravel habitats: assessment and recovery of benthic megafauna on Georges Bank. In: Barnes P, Thomas J (eds) *Benthic habitats and the effects of fishing*, Vol 41. American Fisheries Society, Bethesda, MD., p 325-343
43. Currie DR, et al. (2011) Relative importance of environmental gradients and historical trawling effort in determining the composition and distribution of benthic macro-biota in a large inverse estuary. *Fish Res* 107:184-195

44. Frid CLJ, Clark RA, Hall JA (1999) Long-term changes in the benthos on a heavily fished ground off the NE coast of England. *Mar Ecol Prog Ser* 188:13-20
45. Hiddink JG, Johnson AF, Kingham R, Hinz H (2011) Could our fisheries be more productive? Indirect negative effects of bottom trawl fisheries on fish condition. *J Appl Ecol* 48:1441–1449
46. Hinz H, Prieto V, Kaiser MJ (2009) Trawl disturbance on benthic communities: chronic effects and experimental predictions. *Ecol Appl* 19:761–773
47. Jennings S, et al. (2001) Trawling disturbance can modify benthic production processes. *J Anim Ecol* 70:459-475
48. Jennings S, Nicholson MD, Dinmore TA, Lancaster J (2002) The effect of chronic trawling disturbance on the production of infaunal communities. *Mar Ecol Prog Ser* 243:251-260
49. Kaiser MJ, et al. (2000) Chronic fishing disturbance has changed shelf sea benthic community structure. *J Anim Ecol* 69:494-503
50. Mangano MC, et al. (2014) Infaunal community responses to a gradient of trawling disturbance and a long-term Fishery Exclusion Zone in the Southern Tyrrhenian Sea. *Cont Shelf Res* 76:25-35
51. Poiner I, et al. (1998) The Environmental Effects of Prawn Trawling in the Far Northern Section of the Great Barrier Reef Marine Park: 199-1996 Final Report to the Great Barrier Reef Marine Park Authority and the Fisheries Research & Development Corporation., CSIRO Division of Marine Research, Cleveland
52. Burrridge CY, et al. (2006) A comparison of demersal communities in an area closed to trawling with those in adjacent areas open to trawling: a study in the Great Barrier Reef Marine Park, Australia. *Fish Res* 79:64-74
53. Queirós AM, Hiddink JG, Hinz H, Kaiser MJ (2006) Effects of chronic bottom trawling disturbance on biomass, production and size spectra of invertebrate infauna communities from different habitats. *J Exp Mar Biol Ecol* 335:91-103
54. Reiss H, et al. (2009) Effects of fishing disturbance on benthic communities and secondary production within an intensively fished area. *Mar Ecol Prog Ser* 394:201-213
55. Smith BE, Collie JS, Lengyel NL (2013) Effects of chronic bottom fishing on the benthic epifauna and diets of demersal fishes on northern Georges Bank. *Mar Ecol Prog Ser* 472:199-217
56. Svane I, Hammett Z, Lauer P (2009) Impacts of trawling on benthic macro-fauna and -flora of the Spencer Gulf prawn fishing grounds. *Estuar Coast Shelf Sci* 82:621-631

57. Tillin HM, Hiddink JG, Kaiser MJ, Jennings S (2006) Chronic bottom trawling alters the functional composition of benthic invertebrate communities on a sea basin scale. *Mar Ecol Prog Ser* 318:31-45
58. Veale LO, Hill AS, Hawkins SJ, Brand AR (2000) Effects of long-term physical disturbance by commercial scallop fishing on subtidal epifaunal assemblages and habitats. *Mar Biol* 137:325-337
59. Vergnon R, Blanchard F (2006) Evaluation of trawling disturbance on macrobenthic invertebrate communities in the Bay of Biscay, France: Abundance Biomass Comparison (ABC method). *Aquat Living Resour* 19:219-228
60. Sydow J (1990) Cruise report experiments on the interaction fishing gear (beamtrawl)-benthos with RV Mitra. BEON Rapport 8:1-57
61. Paschen M, Richter U, Köpnick W (2000) TRAPESE - Trawl Penetration in the Seabed. Final Report EU Contract 96-006., University of Rostock
62. Lindeboom HJ, De Groot SJ (1998) IMPACT-II: The effects of different types of fisheries on the North Sea and Irish Sea benthic ecosystems. NIOZ- Report 1998-1/RIVO-DLO Report C003/98. Den Burg, Texel, Netherlands, Netherlands Institute for Sea Research. 404 pp.
63. Bridger J (1972) Some observations on the penetration into the sea bed of tickler chains on a beam trawl. *ICES CM* 7:6
64. Bridger JP, Margetts AR (1971) The effect of a beam trawl on the sea bed. Gear and behaviour committee. *ICES C.M.* 1971/B:8. Report No. Gear and behaviour committee C.M. 1971/B:8
65. Bergman MJN, Hup M (1992) Direct effects of beamtrawling on macrofauna in a sandy sediment in the southern North Sea. *ICES J Mar Sci* 49:5-11
66. Smith C, Rumohr H, Karakassis I, Papadopoulou K-N (2003) Analysing the impact of bottom trawls on sedimentary seabeds with sediment profile imagery. *J Exp Mar Biol Ecol* 285:479-496
67. Schwinghamer P, Guigne J, Siu W (1996) Quantifying the impact of trawling on benthic habitat structure using high resolution acoustics and chaos theory. *Can J Fish Aquat Sci* 53:288-296
68. Freese L, Auster PJ, Heifetz J, Wing BL (1999) Effects of trawling on seafloor habitat and associated invertebrate taxa in the Gulf of Alaska. *Mar Ecol Prog Ser* 182:119-126
69. DEGREE (2010) Development of fishing Gears with Reduced Effects on the Environment (DEGREE). Final Publishable Activity Report-EU Contract SSP8-CT-2004-022576
70. Mengual B, et al. (2016) Influence of bottom trawling on sediment resuspension in the 'Grande-Vasière' area (Bay of Biscay, France). *Ocean Dyn* 66:1181-1207
71. Lucchetti A, Sala A (2012) Impact and performance of Mediterranean fishing gear by side-scan sonar technology. *Can J Fish Aquat Sci* 69:1806-1816

72. Krost P, Bernhard M, Werner F, Hukriede W (1989) Otter trawl tracks in Kiel Bay (Western Baltic) mapped by side-scan sonar. *Meeresforschung* 32:344-353
73. Ivanovic A, Neilson RD, O'Neill FG (2011) Modelling the physical impact of trawl components on the seabed and comparison with sea trials. *Ocean Eng* 38:925–933
74. Humborstad O-B, Nøttestad L, Løkkeborg S, Rapp HT (2004) RoxAnn bottom classification system, sidescan sonar and video-sledge: spatial resolution and their use in assessing trawling impacts. *ICES J Mar Sci* 61:53-63
75. Brylinsky M, Gibson J, Gordon Jr DC (1994) Impacts of flounder trawls on the intertidal habitat and community of the Minas Basin, Bay of Fundy. *Can J Fish Aquat Sci* 51:650-661
76. O'Neill FG, Summerbell K, Breen M (2009) An underwater laser stripe seabed profiler to measure the physical impact of towed gear components on the seabed. *Fish Res* 99:234-238
77. Werner F, et al. (1990) Sedimentological effects of trawl fishery in Kiel Bay(western Baltic). *Meyniana Kiel* 42:123-151
78. Pranovi F, Giovanardi O (1994) The impact of hydraulic dredging for short-necked clams, *Tapes* spp., on an infaunal community in the lagoon of Venice. *Sci Mar* 58:345-353
79. van den Heiligenberg T (1987) Effects of mechanical and manual harvesting of lugworms *Arenicola marina* L. on the benthic fauna of tidal flats in the Dutch Wadden Sea. *Biol Conserv* 39:165-177
80. Clarke S, Tully O (2014) BACI monitoring of effects of hydraulic dredging for cockles on intertidal benthic habitats of Dundalk Bay, Ireland. *J Mar Biol Assoc UK* 94:1451-1464
81. Hall-Spencer JM, Moore PG (2000) Scallop dredging has profound, long-term impacts on maerl habitats. *ICES J Mar Sci* 57:1407-1415
82. Lart W, et al. (2003) Evaluation and improvement of shellfish dredge design and fishing effort in relation to technical conservation measures and environmental impact:[ECODREDGE CT98-4465].
83. Buhl-Mortensen L, et al. (2013) Impacts of fisheries and aquaculture on sediments and benthic fauna: suggestions for new management approaches. *Fisken og Havet* 2:69
84. Caddy JF (1973) Underwater observations on tracks of dredges and trawls and some effects of dredging on a scallop ground. *J Fish Res Board Can* 30:173-180
85. Dale A, Boulcott P, Sherwin T (2011) Sedimentation patterns caused by scallop dredging in a physically dynamic environment. *Mar Pollut Bull* 62:2433-2441
86. de Clerk R, Hovart P (1972) On the effects of tickler chains. *ICES C.M.*1972/B:15, 11 pp.
87. De Groot SJ (1984) The impact of bottom trawling on benthic fauna of the North Sea. *Ocean Management* 9:177-190

88. Depestele J, et al. (2016) Measuring and assessing the physical impact of beam trawling. ICES J Mar Sci 73:i15-i26
89. de Madron XD, et al. (2005) Trawling-induced resuspension and dispersal of muddy sediments and dissolved elements in the Gulf of Lion (NW Mediterranean). Cont Shelf Res 25:2387-2409
90. Gilkinson K, Paulin M, Hurley S, Schwinghamer P (1998) Impact of trawl door scouring on infaunal bivalves: results of a physical trawl door model/dense interaction. J Exp Mar Biol Ecol 224:291-312
91. Houghton R, Williams T, Blacker R (1971) Some effects of double beam trawling. ICES C.M. 1971/B: 5. Gear and Behaviour Committee
92. Laban C, Lindeboom H (1991) Penetration depth of beam trawl gear. BEON Rapport 13:37-52
93. Løkkeborg S (2005) Impacts of trawling and scallop dredging on benthic habitats and communities. Food & Agriculture Org., Rome
94. Journal of Sea Research Fisheries Research Malik MA, Mayer LA (2007) Investigation of seabed fishing impacts on benthic structure using multi-beam sonar, sidescan sonar, and video. ICES J Mar Sci 64:1053-1065
95. O'Neill FG, Summerbell K, Breen M (2008) The Suspension Of Sediment By Scallop Dredges. Fisheries Research Services Internal Report. Report No. No 08/08., FRS Marine Laboratory, Aberdeen, Scotland
96. O'Neill F, et al. (2013) The mobilisation of sediment and benthic infauna by scallop dredges. Mar Environ Res 90:104-112